

Ultrafilters with property (s)

Arnold W. Miller¹

Abstract

A set $X \subseteq 2^\omega$ has property (s) (Marczewski (Szpilrajn)) iff for every perfect set $P \subseteq 2^\omega$ there exists a perfect set $Q \subseteq P$ such that $Q \subseteq X$ or $Q \cap X = \emptyset$. Suppose \mathcal{U} is a nonprincipal ultrafilter on ω . It is not difficult to see that if \mathcal{U} is preserved by Sacks forcing, i.e., it generates an ultrafilter in the generic extension after forcing with the partial order of perfect sets, then \mathcal{U} has property (s) in the ground model. It is known that selective ultrafilters or even P-points are preserved by Sacks forcing. On the other hand (answering a question raised by Hrusak) we show that assuming CH (or more generally MA) there exists an ultrafilter \mathcal{U} with property (s) such that \mathcal{U} does not generate an ultrafilter in any extension which adds a new subset of ω .

It is a well known classical result due to Sierpinski (see [1]) that a nonprincipal ultrafilter \mathcal{U} on ω when considered as a subset of $P(\omega) = 2^\omega$ cannot have the property of Baire or be Lebesgue measurable. Here we identify 2^ω and $P(\omega)$ by identifying a subset of ω with its characteristic function. Another very weak regularity property is property (s) of Marczewski (see Miller [7]). A set of reals $X \subseteq 2^\omega$ has property (s) iff for every perfect set P there exists a subperfect set $Q \subseteq P$ such that either $Q \subseteq X$ or $Q \cap X = \emptyset$. Here by perfect we mean homeomorphic to 2^ω .

It is natural to ask:

Question. (Steprans) Can a nonprincipal ultrafilter \mathcal{U} have property (s)?

If \mathcal{U} is an ultrafilter in a model of set theory V and $W \supseteq V$ is another model of set theory then we say \mathcal{U} generates an ultrafilter in W if for every $z \in P(\omega) \cap W$ there exists $x \in \mathcal{U}$ with $x \subseteq z$ or $x \cap z = \emptyset$. This means that the filter generated by \mathcal{U} (i.e. closing under supersets) is an ultrafilter in W .

We begin with the following result:

¹Thanks to the Fields Institute, Toronto for their support during the time these results were proved and to Juris Steprans for helpful conversations and thanks to Boise State University for support during the time this paper was written.

Mathematics Subject Classification 2000: 03E35; 03E17; 03E50

Theorem 1 *For \mathcal{U} a nonprincipal ultrafilter on ω in V the following are equivalent:*

1. *For some Sack's generic real x over V*

$$V[x] \models \mathcal{U} \text{ generates an ultrafilter.}$$

2. *In V , for every perfect set $P \subseteq P(\omega)$ there exists a perfect set $Q \subseteq P$ and a $z \in \mathcal{U}$ such that either $\forall x \in Q \ z \subseteq x$ or $\forall x \in Q \ z \cap x = \emptyset$.*

3. *For some extension $W \supseteq V$ with a new subset of ω*

$$W \models \mathcal{U} \text{ generates an ultrafilter.}$$

Proof

To see that (3) \rightarrow (2), let P be any perfect set coded in V . Since W contains a new subset of ω there exists $x \in (P \cap W) \setminus V$.

Since \mathcal{U} generates an ultrafilter in W there exists $z \in \mathcal{U}$ so that either $z \subseteq x$ or $z \cap x = \emptyset$. Suppose the first happens. In V consider the set

$$Q = \{y \in P : z \subseteq y\}$$

Note that the new real x is in the closed set Q . It follows that Q must be an uncountable closed set and so it contains a perfect subset. The other case is exactly the same.

One way to see that Q must be uncountable is to note that if (in V) $Q = \{x_n : n < \omega\}$, then the Π_1^1 sentence

$$\forall x \in 2^\omega (x \in Q \text{ iff } \exists n < \omega \ x = x_n)$$

would be true in V and since Π_1^1 sentences are absolute (Mostowski absoluteness, see [5]) true in W . Another way to prove it is to do the standard derivative Cantor argument to the closed set Q removing isolated points and iterating thru the transfinite and noting that each real removed is in V , while the new real is never removed, and hence the kernel of Q is perfect. See Solovay [10], for a similar proof of Mansfield's theorem that a (lightface) Π_2^1 set with a nonconstructible element contains a perfect set.

Now we see that (2) \rightarrow (1). A basic property of Sack's forcing is that for every $y \in 2^\omega \cap M[x]$ in a Sacks extension is either in M or is itself Sacks generic over M (see Sacks [9]). Hence we need only show that if $y \subseteq \omega$ is

Sacks generic over M , then there exists $z \in \mathcal{U}$ with $z \subseteq y$ or $z \cap y = \emptyset$. Recall also that the Sacks real y satisfies that the generic filter G is exactly the set of all perfect sets Q coded in V with $y \in Q$.

Condition (2) says that the set of such Q are dense and hence there exists Q in the generic filter determined by y and $z \in \mathcal{U}$ such that either $z \subseteq u$ for every $u \in Q$ or either $z \cap u = \emptyset$ for every $u \in Q$. But this means that either $z \subseteq y$ or $z \cap y = \emptyset$.

(1) \rightarrow (3) is obvious.

QED

Remark. The above proof also shows that if an ultrafilter is preserved in one Sacks extension, then it is preserved in all Sacks extensions.

Remark. In Baumgartner and Laver [2] it is shown that selective ultrafilters are preserved by Sacks forcing. In Miller [6] it is shown that P -points are preserved by superperfect set forcing (and hence by Sacks forcing also).

We say that an ultrafilter \mathcal{U} is preserved by Sacks forcing iff for some (equivalently all) Sacks generic reals x that \mathcal{U} generates an ultrafilter in $V[x]$. Recall that $\mathcal{U} \times \mathcal{V}$ is the ultrafilter on $\omega \times \omega$ defined by

$$A \in \mathcal{U} \times \mathcal{V} \text{ iff } \{n : \{m : (n, m) \in A\} \in \mathcal{U}\} \in \mathcal{V}$$

If \mathcal{U} and \mathcal{V} are nonprinciple ultrafilters, then $\mathcal{U} \times \mathcal{V}$ is not a P-point. Also recall that $\mathcal{U} \leq_{RK} \mathcal{V}$ (Rudin-Keisler) iff there exists $f \in \omega^\omega$ such that for every $X \subseteq \omega$

$$X \in \mathcal{U} \text{ iff } f^{-1}(X) \in \mathcal{V}$$

Proposition 2 *If \mathcal{U} and \mathcal{V} are preserved by Sacks forcing, then so is $\mathcal{U} \times \mathcal{V}$. If $\mathcal{U} \leq_{RK} \mathcal{V}$ and \mathcal{V} is preserved by Sacks forcing, then so is \mathcal{U} .*

Proof

Suppose $A \subseteq \omega \times \omega$ and $A \in V[x]$. For each $n < \omega$ let $A_n = \{m : (n, m) \in A\}$. Since \mathcal{U} is preserved there exists $B_n \in \mathcal{U}$ with $B_n \subseteq A_n$ or $B_n \cap A_n = \emptyset$. By the preservation of \mathcal{V} there exists $C \in \mathcal{V}$ such that either $B_n \subseteq A_n$ for all $n \in C$ or $B_n \cap A_n = \emptyset$ for all $n \in C$. By the Sacks property there exists $(\mathfrak{b}_n \in [\mathcal{U}]^{2^n} : n < \omega) \in V$ such that $B_n \in \mathfrak{b}_n$ for every n . Let $B_n^0 = \cap \mathfrak{b}_n$. Then

$$\cup_{n \in C} \{n\} \times B_n^0 \subseteq A \text{ or } \cup_{n \in C} \{n\} \times B_n^0 \cap A = \emptyset$$

Suppose $\mathcal{U} \leq_{RK} \mathcal{V}$ via f . If $A \subseteq \omega$, then since \mathcal{V} is preserved, there exists $B \in \mathcal{V}$ such that either $B \subseteq f^{-1}(A)$ or $B \subseteq f^{-1}(\overline{A})$. but then $f(B) \subseteq A$ or $f(B) \subseteq \overline{A}$ and since $f(B) \in \mathcal{U}$ we are done.
QED

Remark. The Rudin-Keisler result is generally true, but the product result depends on the bounding property. For example, if \mathcal{U} is a P-point, then \mathcal{U} is preserved in the superperfect extension, but $\mathcal{U} \times \mathcal{U}$ is not.

It is clear that property (2) of Theorem 1 implies that any ultrafilter which is preserved by Sacks forcing has property (s). But what about the converse? The main result of this paper is that the reverse implication is false. This answers a question raised by Hrusak.

Theorem 3 *Suppose the CH is true or even just that the real line cannot be covered by fewer than continuum many meager sets. Then there exists an ultrafilter \mathcal{U} on ω which has property (s) but is not preserved by Sacks forcing.*

Proof

We give the proof in the case of the continuum hypothesis and indicate how to do it under the more general hypothesis.

Let $\mathcal{I} \subseteq [\omega]^\omega$ be an independent perfect family. Independent means that for every m, n and distinct $x_1, \dots, x_m, y_1, \dots, y_n \in \mathcal{I}$ the set

$$x_1 \cap \dots \cap x_m \cap \overline{y}_1 \cap \dots \cap \overline{y}_n \text{ is infinite.}$$

where \overline{y} means the complement of y in ω . We claim that the following family

$$\mathcal{I} \cup \{\overline{z} : \exists^\infty x \in \mathcal{I} \ z \subseteq^* x\}$$

has the finite intersection property. (\exists^∞ means there exists infinitely many). To see this suppose that $x_1, \dots, x_m \in \mathcal{I}$ and z_1, \dots, z_n and $\exists^\infty x \in \mathcal{I} \ z_i \subseteq^* x$ for each i . Then we can choose $y_i \in \mathcal{I}$ distinct from each other and the x 's so that each $z_i \subseteq^* y_i$. But since $\overline{y}_i \subseteq \overline{z}_i$ we have that

$$x_1 \cap \dots \cap x_m \cap \overline{y}_1 \cap \dots \cap \overline{y}_n \subseteq^* x_1 \cap \dots \cap x_m \cap \overline{z}_1 \cap \dots \cap \overline{z}_n$$

By independence the set on the left is infinite and hence so is the set on the right. Thus this family has the finite intersection property.

Now let \mathcal{F}_0 be the filter generated by $\mathcal{I} \cup \{\bar{z} : \exists^\infty x \in \mathcal{I} \ z \subseteq^* x\}$.

Note that if $\mathcal{U} \supseteq \mathcal{F}_0$ is any ultrafilter then it cannot be preserved by Sacks forcing. This is because \mathcal{I} is a perfect subset of \mathcal{U} , however there is no $z \in \mathcal{U}$ with $z \subseteq x$ for all $x \in \mathcal{I}$ or even infinitely many $x \in \mathcal{I}$ or else $\bar{z} \in \mathcal{F}_0 \subseteq \mathcal{U}$, hence Theorem 1 (2) fails.

Note that since \mathcal{I} was perfect the filter \mathcal{F}_0 is a Σ_1^1 subset of $P(\omega)$.

Lemma 4 *Suppose that $P \subseteq [\omega]^\omega$ is a perfect set and \mathcal{F} is a Σ_1^1 filter extending the cofinite filter on ω . Then there exists a perfect $Q \subseteq P$ such that either*

1. $\mathcal{F} \cup Q$ has the finite intersection property or
2. there exists $z \subseteq \omega$ so that $\mathcal{F} \cup \{z\}$ has the finite intersection property and for every $x \in Q$ we have that there exists $y \in \mathcal{F}$ with $x \cap y \cap z = \emptyset$.

Proof

The strategy is try to do a fusion argument to get case (1). If it every fails, then stop and get case (2).

Claim. Suppose $(Q_i : i < n)$ are disjoint perfect subsets of $[\omega]^\omega$. Then either there exists $(Q'_i \subseteq Q_i : i < n)$ perfect so that for every $(x_i \in Q'_i : i < n)$ and $y \in \mathcal{F}$ we have that

$$|y \cap x_0 \cap x_1 \cap \dots \cap x_{n-1}| = \omega$$

or there exists $z \subseteq \omega$, $k < n$, and $Q \subseteq Q_k$ perfect so that $\mathcal{F} \cup \{z\}$ has the finite intersection property and for every $x \in Q$ we have that there exists $y \in \mathcal{F}$ with $x \cap y \cap z = \emptyset$.

Proof

Consider

$$A_k = \{(x_i \in [\omega]^\omega : i < k) : \exists y \in \mathcal{F} \ |y \cap x_0 \cap x_1 \cap \dots \cap x_{k-1}| < \omega\}$$

Since \mathcal{F} is Σ_1^1 it is easy to see that each A_k is a Σ_1^1 set and hence has the property of Baire relative to the product $\prod_{i < k} Q_i$. By Mycielski [8] (see also Blass [3]) there exists perfect sets $(Q_i^* \subseteq Q_i : i < n)$ so that for every $k \leq n$ either

$$\prod_{i < k} Q_i^* \cap A_k = \emptyset \text{ or } \prod_{i < k} Q_i^* \subseteq A_k.$$

If the first case happens for $k = n$, then we let $Q'_i = Q_i^*$ and the claim is proved. If the second case happens choose k minimal for which it happens. This means we have that

1. for all $(x_i : i < k - 1) \in \prod_{i < k-1} Q_i^*$ and $y \in \mathcal{F}$ we have

$$|y \cap x_0 \cap x_1 \cap \dots \cap x_{k-2}| = \omega$$

and

2. for all $(x_i : i < k) \in \prod_{i < k} Q_i^*$ there exists $y \in \mathcal{F}$ such that

$$|y \cap x_0 \cap x_1 \cap \dots \cap x_{k-1}| = \emptyset$$

In this case let $(x_i : i < k - 1) \in \prod_{i < k-1} Q_i^*$ be arbitrary and put

$$z = x_0 \cap x_1 \cap \dots \cap x_{k-2} \text{ and } Q = Q_k^* \subseteq Q_k.$$

This proves the Claim.

QED

It is now an easy fusion argument to finish proving the Lemma from the Claim.

QED

Now we construct our ultrafilter proving the theorem under the assumption of CH. We let $(P_\alpha : \alpha < \omega)$ list all perfect subsets of 2^ω . We construct an increasing sequence \mathcal{F}_α for $\alpha < \omega_1$ of Σ_1^1 filters as follows.

Let \mathcal{F}_0 the filter generated by

$$\mathcal{I} \cup \{\bar{z} : \exists^\infty x \in \mathcal{I} \ z \subseteq^* x\}$$

At limit ordinals α we let \mathcal{F}_α be the union $\cup_{\beta < \alpha} \mathcal{F}_\beta$ and note that it is a Σ_1^1 filter. At successor stages $\alpha + 1$ we apply the Lemma to P_α and \mathcal{F}_α . In the first case we find a perfect set $Q \subseteq P_\alpha$ such that $\mathcal{F}_\alpha \cup Q$ has the finite intersection property. In this case we let $\mathcal{F}_{\alpha+1}$ be the filter generated by $\mathcal{F}_\alpha \cup Q$ and note that it is Σ_1^1 . In the second case we find a perfect set $Q \subseteq P_\alpha$ and $z \subseteq \omega$ so that $\mathcal{F}_\alpha \cup \{z\}$ has the finite intersection property and for every $x \in Q$ we have that there exists $y \in \mathcal{F}_\alpha$ with $x \cap y \cap z = \emptyset$. Here we let $\mathcal{F}_{\alpha+1}$ be the filter generated by $\mathcal{F}_\alpha \cup \{z\}$ and note that for every ultrafilter $\mathcal{U} \supseteq \mathcal{F}_{\alpha+1}$ that $\mathcal{U} \cap Q = \emptyset$, because we have put $\{\bar{x} : x \in Q\} \subseteq \mathcal{F}_{\alpha+1}$. this ends the proof under CH.

Now we see how do this construction under the weaker hypothesis that the real line cannot be covered by fewer than continuum many meager sets. We construct an increasing sequence $(\mathcal{F}_\alpha : \alpha < \mathfrak{c})$ of filters such that each \mathcal{F}_α is the union of $\leq |\alpha|$ Σ_1^1 sets. In order to prove the corresponding Claim and Lemma we note that the following is true.

Claim Suppose the real line cannot be covered by κ many meager sets, $(Q_k : k < n)$ are perfect, and for each $k \leq n$ we have $A_k \subseteq \prod_{i < k} [\omega]^\omega$ which is the union of $\leq \kappa$ many Σ_1^1 sets. Then there exists $(Q_i^* \subseteq Q_i : i < n)$ perfect so that for every $k \leq n$ either

$$\prod_{i < k} Q_i^* \subseteq A_k \text{ or } \prod_{i < k} Q_i^* \cap A_k = \emptyset.$$

Proof

Construct $(Q_i^j : i < n)$ perfect by induction so that

1. $Q_i^0 = Q_i$ all $i < n$,
2. $Q_i^{j+1} \subseteq Q_i^j$ all $i < n$,
3. $\prod_{i < j} Q_i^j \subseteq A_j$ or $\prod_{i < j} Q_i^j \cap A_j = \emptyset$.

Given $(Q_i^j : i < n)$ and A_{j+1} the union of κ many Σ_1^1 sets, say $\cup \{B_\alpha : \alpha < \kappa\}$ there are two cases.

Case 1. For some $\alpha < \kappa$ the set $B_\alpha \cap \prod_{i < j+1} Q_i^j$ is not meager.

In this case it must be comeager in some relative interval $\prod_{i < j+1} Q_i^j \cap \prod_{i < j+1} [s_i]$. And now we can find $Q_i^{j+1} \subseteq Q_i^j \cap [s_i]$ such that

$$\prod_{i < j+1} Q_i^{j+1} \subseteq B_\alpha \subseteq A_{j+1}$$

Case 2. Each B_α is meager in $\prod_{i < j+1} Q_i^j$.

In this case we use the covering of category hypothesis in the form of Martin's axiom for countable posets. For $p \subseteq \omega^{<\omega}$ a finite subtree, define s a terminal node of p ($s \in \text{term}(p)$) iff $s \in p$ and for every $t \in p$ if $s \subseteq t$ then $s = t$. For p, q finite subtrees of $2^{<\omega}$ we define $p \supseteq_e q$ (end extension) iff $p \supseteq q$ and every new node of p extends a terminal node of q . Define $T_i = \{s \in 2^{<\omega} : [s] \cap Q_i \neq \emptyset\}$.

Consider the partial order \mathbb{P} consisting of finite approximations to products of perfect trees below the Q_i :

$$\mathbb{P} = \{(p_i : i < n) : p_i \text{ is a finite subtree of } T_i, i < n\}$$

and $p \leq q$ iff $p_i \supseteq_e q_i$ all $i < n$. Our assumption about the covering of the real by meager sets is equivalent to $\text{MA}_\kappa(\text{ctble})$, i.e., for every countable poset \mathbb{P} and any family $(D_\alpha : \alpha < \kappa)$ of dense subsets of \mathbb{P} there exists a \mathbb{P} -filter G such that $G \cap D_\alpha \neq \emptyset$ for all $\alpha < \kappa$. Note that for any $k < n$ and $C \subseteq \prod_{i < k} Q_i$ which is nowhere dense the set

$$D_C = \{p \in \mathbb{P} : \forall i < k \ \forall (s_i \in \text{term}(p_i) : i < k) \ C \cap \prod_{i < k} [s_i] = \emptyset\}$$

is dense in \mathbb{P} similarly, for any $m < \omega$ the following sets are dense:

$$D_m = \{p \in \mathbb{P} : \forall i < n \forall s \in \text{term}(p_i) \ |s| > m\}$$

$$D_m^* = \{p \in \mathbb{P} : \forall i < n \forall s \in p_i \ |s| = m \rightarrow \exists t \ t \supseteq s \text{ and } t0, t1 \in p_i\}$$

So a sufficiently generic filter produces a sequence $(T'_i \subseteq T_i : i < n)$ of perfect subtrees such that letting $Q'_i = [T'_i]$ we have the property that $\prod_{i < n} Q'_i \cap B_\alpha = \emptyset$ for all $\alpha < \kappa$.
QED

Question 5 *Can we prove in ZFC that there exists a nonprincipal ultrafilter with property (s)?*

Remark. It is easy to construct a nonprincipal ultrafilter which fails to have property (s). Start with a perfect independent family $I \subseteq P(\omega)$. Choose

$$\{X_\alpha : \alpha < \mathfrak{c}\} \cup \{Y_\alpha : \alpha < \mathfrak{c}\} \subseteq I$$

distinct so that for every perfect $Q \subseteq I$ there exists α with X_α and Y_α both in Q . Then any ultrafilter

$$\mathcal{U} \supseteq \{X_\alpha : \alpha < \mathfrak{c}\} \cup \{\overline{Y}_\alpha : \alpha < \mathfrak{c}\}$$

will fail to have property (s).

Question 6 *Can we prove in ZFC that there exists a nonprinciple ultrafilter with which is preserved by Sacks forcing?*

Note Shelah, see [1], has shown it is consistent that there are no non-principle P-points. See Brendle [4] for a plethora of ultrafilters weaker than P-points such as Baumgartner’s nowhere dense ultrafilters.

Question 7 *Suppose \mathcal{U} and \mathcal{V} are nonprinciple ultrafilters in V which generate ultrafilters \mathcal{U}^* and \mathcal{V}^* in $W \supseteq V$. If $\mathcal{U}^* \leq_{RK} \mathcal{V}^*$ holds in W , must $\mathcal{U} \leq_{RK} \mathcal{V}$ be true in V ?*

Question 8 *Suppose $\mathcal{U} \in V$ is generates an ultrafilter in $V[x]$ for some (equivalently all) Sacks reals x over V . Suppose x is a Sacks real over V and y is a Sacks real over $V[x]$. Must \mathcal{U} generate an ultrafilter in $V[x, y]$?*

Question 9 *Suppose $V \subseteq W$ and*

$$V \models \mathcal{U} \text{ has property } (s)$$

does

$$W \models \{A \subseteq \omega : \exists B \in \mathcal{U} B \subseteq A\} = \mathcal{U}^* \text{ has property } (s) \text{ ?}$$

Question 10 *Is Proposition 2 true for property (s) in place of “preserved by Sacks forcing”?*

References

- [1] Bartoszyński, Tomek; Judah, Haim; **Set theory. On the structure of the real line.** A K Peters, Ltd., Wellesley, MA, 1995. (Ultrafilters can’t be measurable or have property of Baire.)
- [2] Baumgartner, James E.; Laver, Richard; Iterated perfect-set forcing. *Ann. Math. Logic* 17 (1979), no. 3, 271–288.
- [3] Blass, Andreas; A partition theorem for perfect sets. *Proc. Amer. Math. Soc.* 82 (1981), no. 2, 271–277.
- [4] Brendle, Jorg; Between P -points and nowhere dense ultrafilters. *Israel J. Math.* 113 (1999), 205–230.

- [5] Kechris, Alexander S.; **Classical descriptive set theory.** Graduate Texts in Mathematics, 156. Springer-Verlag, New York, 1995.
- [6] Miller, Arnold W.; Rational perfect set forcing. Axiomatic set theory (Boulder, Colo., 1983), 143–159, Contemp. Math., 31, Amer. Math. Soc., Providence, RI, 1984.
- [7] Miller, Arnold W.; Special subsets of the real line. Handbook of set-theoretic topology, 201–233, North-Holland, Amsterdam, 1984.
- [8] Mycielski, Jan; Independent sets in topological algebras. Fund. Math. 55 1964 139–147.
- [9] Sacks, Gerald E.; Forcing with perfect closed sets. 1971 Axiomatic Set Theory (Proc. Sympos. Pure Math., Vol. XIII, Part I, Univ. California, Los Angeles, Calif., 1967) pp. 331–355 Amer. Math. Soc., Providence, R.I.
- [10] Solovay, Robert M.; On the cardinality of \sum_2^1 sets of reals. 1969 Foundations of Mathematics (Symposium Commemorating Kurt Godel, Columbus, Ohio, 1966) pp. 58–73 Springer, New York.

Arnold W. Miller
 miller@math.wisc.edu
<http://www.math.wisc.edu/~miller>
 University of Wisconsin-Madison
 Department of Mathematics, Van Vleck Hall
 480 Lincoln Drive
 Madison, Wisconsin 53706-1388